



MATRIXYL®3000



Pal-GHK and Pal-GQPR

Function:
Anti-wrinkle.

Definition:
Association of 2 palmitoylated matrikines: Pal-Gly-His-Lys and Pal-Gly-Glu-Pro-Arg.

Properties:
Matrixyl®3000 supports the activation of the cutaneous repair process, and in particular at the level of the fragile and UV-damage prone papillary dermis. Matrixyl®3000 promotes wrinkle smoothing and improves tone and elasticity.

Characteristics:
Matrikines are messenger peptides capable of regulating cell activities by interacting with their specific receptors. They activate certain genes involved in the process of extracellular matrix renewal and cell proliferation. These mechanisms become weaker and weaker with age.

INCI Name:
(Check PCPC on-line dictionary for latest INCI name)
Glycerin – Aqua (Water) – Butylene Glycol – Carbomer – Polysorbate 20 – Palmitoyl Oligopeptide – Palmitoyl Tetrapeptide-7

Applications:
Anti-wrinkle products.


Formulation:
Water soluble.

Recommended use level:
3%

Patent:
WO 2005/048968



NEW TEST reveals age gain by **2** years in just **1** month



www.sederma.fr
E-mail: sederma@sederma.fr
Copyright© 2000-2011
Sederma. All rights reserved.

Member of Croda International Plc


In vitro tests

● SYNTHESIS OF MATRIX MACROMOLECULES

Study of the stimulation of synthesis of extracellular matrix components by fibroblasts incubated for 72 hours with Matrixyl®3000 (1, 3, 5%).

● STIMULATION OF GENE EXPRESSION

Study of the regulation of dermal and epidermal genes by matrikines from Matrixyl®3000, using DNA-Array technique on reconstructed epidermis and fibroblast culture.



Matrixyl®3000 can boost the synthesis of extracellular matrix macromolecules. Its matrikines showed a complementary activation profile of genes involved in the mechanism of cutaneous restructuring.


In vivo tests

REPAIR OF THE PAPILLARY DERMIS

28 female volunteers aged from 51 to 72 years, mean age 59. Twice daily application of a cream containing 3% of Matrixyl®3000 for 2 months to one half of the face and the forearm (inner and UV-exposed outer forearm) against placebo.

● ANALYSIS OF THE SUBEPIDERMAL LOW ECHOGENICITY BAND


after 2 months	Inner forearm	Outer forearm
THICKNESS	-9.8%/T0 up to -23%, p<0.01	-9.8%/T0 up to -33%, p<0.01
	93% volunteers	86% volunteers
	-11%/placebo, p<0.01	-14.4%/placebo, p<0.01
DENSITY	+11.4%/T0 up to +44%, p<0.01	+11.5%/T0 up to +45%, p<0.01
	68% volunteers	82% volunteers
	+15.2%/placebo, p<0.01	+15.1%/placebo, p<0.01
AGE GAIN	-3.8 years	-5.5 years



Significant improvement of the SLEB characteristics in just one month (thickness: -5.5%/placebo; density: +7.8%/placebo, on the inner arm) and confirmed after two months. The ageing process is slowed down by 1.8 years in just one month.

● ANALYSIS OF THE FIBRE NETWORK

Measurements of the fibre defragmentation by confocal laser microscopy (Vivascope) on the face, next to the external eye corner.



Matrixyl®3000 helps reduce the fibre fragmentation and notably supports the reconstruction of the papillary dermal fibre network.

ANTI-WRINKLE EFFICACY, SKIN TONE AND ELASTICITY

23 female volunteers aged from 42 to 67 years / Twice daily application on one half of the face of a cream containing 3% of Matrixyl®3000 against placebo, for 2 months. Assessment of the anti-wrinkle efficacy by profilometry, cutometry and photography compared to T0.

Compared to T0 (%)	Matrixyl®3000	Placebo
Surface occupied by deep wrinkles	- 39.4**	4.3ns
Main wrinkle density	- 32.9**	- 9.6ns
Main wrinkle average depth	- 19.9**	- 3.2ns
Main wrinkle average volume	- 23.3**	- 8.7*
Roughness	- 16.0**	1.4ns
Complexity (Lifting effect)	- 16.2**	4.2ns
Elasticity	+5.5*	4.1ns
Skin tone	+15.5**	6.5ns

ns : non significant *significant/T0 (p<0.05) **significant/T0 (p<0.01)



The repairing effect of Matrixyl®3000 promotes the visible quality of the skin by decreasing the appearance of wrinkles and improving tone and elasticity.